

On the Topological Origin of Entanglement in Ising Spin Glasses

V. V. Sreedhar*

*Department of Physics
Indian Institute of Technology
Kanpur, 208016 India*

The origin of thermal and quantum entanglement in a class of three-dimensional spin models, at low momenta, is traced to purely topological reasons. The establishment of the result is facilitated by the gauge principle which, when used in conjunction with the duality mapping of the spin models, enables us to recast them as lattice Chern-Simons gauge theories. The thermal and quantum entanglement measures are expressed in terms of the expectation values of Wilson lines, loops, and their generalisations. For continuous spins, these are known to yield the topological invariants of knots and links. For Ising-like models, they are expressible in terms of the topological invariants of three-manifolds obtained from finite group cohomology – the so-called Dijkgraaf-Witten invariants.

PACS numbers: 03.65.Ud, 11.15.Ha, 11.15.-q, 02.10.Kn

The history of modern physics is replete with stories of the progress that followed every time the notion of instantaneous action at a distance between two objects was sacrificed in favour of the concept of an interaction between the participants. With an appropriate choice of gauge fields acting as mediators of the interaction, we can account for most of what is known about the fundamental forces of nature. Several examples readily come to mind *viz.* Maxwell's theory of electromagnetism, the standard model of the electroweak forces, quantum chromodynamics – the theory of strong, nuclear forces – and Einstein's general theory of relativity for gravitational forces [1]. Drawing inspiration from these examples, we shall use the idea of gauge invariance as a *tour de force* to gain insight into the origin of entanglement in Ising spin systems.

The need to understand the origin of quantum entanglement arises from the recognition that it is the theoretical bedrock that supports the emerging revolution in storing, processing, and retrieving information [2]. Most of the work on the physics of quantum entanglement is – in view of its aforementioned technological import – focussed on finding ways to quantify it, in developing sophisticated algorithms to handle computational complexity, in grappling with the issue of decoherence, and last but not least, in producing experimental devices which shape the promised revolution. Under the circumstances, very little time is invested in finding an answer to a fundamental question: What is the origin of entanglement in quantum systems? We will show that the gauge principle once again holds the key to unlock this, yet another, mystery of nature.

In view of the fact that entanglement is a purely quantum mechanical feature, unlike the fundamental forces of nature alluded to earlier, it may seem that the gauge

principle is inapplicable in this context. Such apprehensions are unfounded because it is by now well-known that purely quantum features with no classical analogues are often associated with non-trivial topological aspects of the theory. The simplest example which illustrates this intimate relation is that of a free nonrelativistic point particle moving on a circle. A total derivative term added to the Lagrangian of this model leaves the classical mechanics of the particle unchanged, but leads to different, inequivalent quantizations of the particle due to the topology of the circle [3]. Another example concerns identical particles which acquire anyonic quantum statistics in a plane, due to the non-trivial topology of the configuration space [4]. The important role played by topological objects like monopoles and vortices in producing confinement, and instantons in producing anomalies, in Yang-Mills gauge theories, is also well documented [5]. The reasons for the quantization of charge in electrodynamics, flux in superconductivity and conductance in the Hall effect can all be traced to their topological antecedents [6]. It should be noted that for each of these examples, there is no classical counterpart; the physical effects have a topological origin, and are most elegantly captured by the gauge principle. We then conjecture that quantum entanglement is also purely topological in origin and substantiate this claim by using the gauge principle, in the concrete example of the three-dimensional Ising model.

We mention that an intriguing analogy between quantum entanglement and classical topology was first publicised by Arvind [7], following his serendipitous discovery of the similarity between the entanglement properties of the Greenberger-Horne-Zeilinger state [8] and the curious linking properties of Borromean and Hopf rings. The analogy was further developed by Kauffman and Lomonaco [9] who examined the parallels between entanglement of states in quantum mechanics on the one hand, and the strings of a braid on the other. The hope that the breakthrough in finding new topological invari-

*sreedhar@iitk.ac.in

ants of knots and links [10] may be used to characterize/classify entangled quantum states through the above analogy, gave rise to some initial excitement. The limitations of such a hope, as delineated in [7], were primarily because the much-needed rule to associate a closed loop to a quantum state, and the snipping of a loop to a measurement, could not be formalised. In the present work we would like to approach this problem from Heisenberg's point of view, and deal directly with the physically observable quantities. This approach enables us to circumvent the need to define the topological equivalents of spins and measurements. We thus overcome the temptation to look for parallels between quantum entanglement and classical topology. Instead, we search for the intrinsic topological content of entanglement in a prototypical spin system namely, the three-dimensional Ising model. In the process, we discover unambiguous relations between the usual measures of entanglement [11] and the expectation values of gauge-invariant observables in a Z_2 lattice Chern-Simons gauge theory. As is well-known, the latter are just the observables used to calculate topological invariants of knots and links in topological field theories with continuous groups [12]. For the case of finite groups, the appropriate topological invariants, obtained by using finite group cohomology, are called the Dijkgraaf-Witten invariants [13].

The Ising model is, as usual, defined by the Hamiltonian $H = -J \sum_{\langle ij \rangle} S_i S_j$, where $J > 0$ and $S_i = \pm 1$. Here i, j label the sites of a three-dimensional cubical lattice and the $\langle \rangle$ parantheses indicate that the summation is over different, but nearest neighbour sites. The positivity of J implies that the ferromagnetic state minimises the energy. For the opposite sign of J , the antiferromagnetic state would be favoured. As already motivated, we proceed to replace the above Hamiltonian which couples spins at different sites, by

$$H_U = -J \sum_{\langle ij \rangle} S_i U_{ij} S_j \quad (1)$$

in which the interaction between spatially separated spins is mediated by the gauge field U . The U_{ij} live on the links connecting sites i and j , are Z_2 -valued, and hence equal to ± 1 . The gauged Ising model presented above accommodates either a ferromagnetic or an antiferromagnetic bond between various nearest neighbour sites [14]. Such models play a crucial role in understanding the behaviour of disordered systems; in which subject they are referred to, and extensively studied, as *spin glasses* [15]. In this letter, however, a spin glass is merely an expedient to replace the action at a distance (of the order of the lattice spacing) between Ising spins by an interaction mediated by the gauge fields U_{ij} .

We define the partition function for the above model in the usual way as the trace over the Gibbs' measure *i.e.* $Z = \text{Tr } e^{-\beta H_U}$ where $\beta = 1/k_B T$, k_B being the

Boltzmann constant, and T , the temperature. In the first step, called *quenching* in the language of spin glasses, we sum over all the spin degrees of freedom to get [16]

$$Z = A(J) \sum_{U,V} e^{-S_V - S_{CS}} \quad (2)$$

where

$$S_V = -\tilde{J} \beta \sum_{\square} \prod_{\square} V_{ij} \quad (3)$$

and

$$S_{CS} = \beta \sum_{\langle ij \rangle} i \frac{\pi}{4} (1 - \prod_{\square} V)(1 - U) \quad (4)$$

The constants A and $\tilde{J}$ are defined in terms of the strength of the magnetic interaction J , as follows:

$$A(J) = \cosh J \quad \text{and} \quad \tanh \tilde{J} = e^{-2J} \quad (5)$$

The V_{ij} , like the U_{ij} , are Z_2 -valued fields, but live on the links of the dual lattice obtained by a rigid translation of each site on the primary lattice by half a lattice spacing uniformly along each of the coordinate axes. The product of the dual gauge variables V_{ij} around an elementary plaquette on the dual lattice is indicated by the $\square$ under the product symbol. When the $\square$ appears under the summation symbol, it is an instruction to sum over all such elementary plaquettes in the dual lattice. The S_V term is readily recognised as the standard Wilsonian action for the gauge field V . The S_{CS} term is physically a measure of the flux passing through a dual plaquette perpendicular to a given link U on the primary lattice. It is a Chern-Simons action on the lattice. Two crucial steps in arriving at the above result consist in an expansion of Z in the characters of the Z_2 group, and the introduction of the dual lattice variables. The details can be found in [16]. The generalisation to other finite abelian groups Z_p follows along the same lines [17], as does the limiting case $p \rightarrow \infty$. The latter corresponds to the familiar lattice $U(1)$ Chern-Simons theory [18].

At this stage, a few general observations regarding the origin of the various terms in (2) are in order. First, from the continuum perspective, for the case of a continuous group, this result is easily anticipated. It is well-known from [19] that a derivative expansion of the fermionic determinant in this case, obtained by integrating out the fermions coupled to a background gauge field in three-dimensional spacetime, produces at the lowest two orders, the Chern-Simons and Maxwell terms. The result in (2-5) is a lattice realisation of the above celebrated continuum result. Second, from a purely lattice point of view, it is well-known that the three-dimensional Ising model is equivalent to a Z_2 lattice gauge theory on the dual lattice [20]. The Ising spins with nearest neighbour

interactions on the primary lattice have thus been traded for the V -fields with a Wilson action that appears in (3). In the duality transformation, the U fields introduced by the gauge principle on the primary lattice are mere spectators and hence we arrive at a set of two Z_2 -valued fields. Moreover, since there is an inextricable linkage between the primary and dual lattices, each link on the primary lattice pierces a plaquette of the dual lattice (and *vice versa*), and the Chern-Simons action is precisely a measure of this flux. Third, the noticeable difference between the continuum and lattice realisations of what is essentially the same result is reminiscent of the lattice fermion doubling problem and has been discussed before in attempts to discretise Chern-Simons gauge theories [21]. It may be recalled for the sake of completeness that, the mathematical reason for the doubling is that the Hodge star operator appearing in the Chern-Simons term couples cochains of a simplicial decomposition with the cochains of the dual decomposition. From a physical point of view, the Chern-Simons term, owing to the fact that it has no independent gauge-invariant dynamics, couples matter fields to the magnetic flux. In the present case, the Ising spins residing on the sites of the primary lattice are the matter fields. The magnetic flux, in lattice gauge theory, is defined by the gauge fields on plaquettes. There is no natural way of coupling these two objects without doing violence to the structure of the lattice. This very fact was used by Kantor and Susskind for one of the early models for anyons [22].

Finally, we mention that models of topological quantum computation using anyons were proposed in [23] and experiments with flux-qubits [24] are beginning to realise some of these ideas. Quite independently, remarkable progress has been achieved by using statistical mechanical techniques to study spin glasses in theories of information processing [25] following the proposal of [26]. Interestingly, equation (2) establishes a connection between the above two seemingly different approaches to the subject of quantum information. More importantly, since the conceptual roots of the former approach lie in topology, while those of the latter lie in gauge invariance, it reinforces the idea anticipated at the beginning of this letter.

To proceed further, we note that although equation (4) does not treat U and V on the same footing, the exponential of $-S_{CS}$ which appears in equation (2) is invariant under an exchange of U and V [16]. However, equation (2) itself is lopsided because it does not have a Wilson term associated with the U field. This suggests that in the next step, called *configurational averaging*, where the quenched gauge degrees of freedom are summed over subject to a given distribution, we choose the weights such that the $U \leftrightarrow V$ symmetry is restored. This is tantamount to using $H_U + S_U$ where $S_U = -K \sum_{\square} \prod_{\square} U_{ij}$ instead of H_U . It is clear from the context that, in this case, the plaquettes under consideration belong to the

primary lattice. If we now introduce the two-component vector $\Omega = (U, V)$ and the matrices

$$M = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad N = \begin{pmatrix} K & 0 \\ 0 & \tilde{J} \end{pmatrix} \quad (6)$$

the partition function can be rewritten in the neat form

$$Z = \text{Tr} e^{-\beta(H_U + K S_U)} = A(J) \sum_{\Omega} e^{-S} \quad (7)$$

where

$$S = \frac{\beta}{2} \sum_{\langle ij \rangle} i \frac{\pi}{4} (1 - \prod_{\square} \Omega) M (1 - \Omega) - \beta N \sum_{\square} \prod_{\square} \Omega_{ij} \quad (8)$$

The $\langle ij \rangle$ in the above equation refers to, as before, nearest neighbour sites, but these could now be either on the primary or dual lattice. Next we take the infra-red (low momentum) limit and drop the N -term to get,

$$S = \frac{\beta}{2} \sum_{\langle ij \rangle} i \frac{\pi}{4} (1 - \prod_{\square} \Omega) M (1 - \Omega) \quad (9)$$

This is recognised as the lattice version of the topological Chern-Simons theory encountered in the studies of the double layer quantum Hall effect [27], and sometimes also referred to as the BF-theory [28].

We are now in a position to examine the entanglement properties of Ising spins in the above system. The object of central interest in studying entanglement is the (reduced) density matrix. Once it is known, a relatively straightforward calculation yields the von Neumann entropy S of the system through the standard formula $S = -\text{Tr} \rho \ln \rho$. The single particle reduced density matrix ρ_i , obtained by tracing over all the spins except the i -th spin, can be expanded as $\rho_i = \frac{1}{2} \sum_{\alpha=0}^3 c_{\alpha} \sigma_i^{\alpha}$ where $\sigma^0 = \mathbf{1}$ and $\sigma_i^{\alpha \neq 0}$ are the Pauli matrices at site i . The requirement that $\text{Tr} \rho_i = 1$, the reality of H , and the global spin-flip symmetry of the Hamiltonian imply that apart from c_0 which is unity, c_1 is the only non-vanishing expectation value. A general expression for the coefficients c_{α} reads

$$c_{\alpha} = \frac{1}{Z} \langle \sigma_i^{\alpha} \mathcal{P} \prod_{\Gamma(i,j=\infty)} U_{ij} \rangle \quad (10)$$

The correlation function has been modified by the path-ordered insertion of a string Γ of links connecting the point i to ∞ making it path-dependent; the modification ensures that the density matrix is gauge-invariant. In a similar fashion, we can obtain the two-particle reduced density matrix by expanding it in terms of the tensor product of Pauli matrices at the two sites under consideration: $\rho_{ij} = \frac{1}{4} \sum_{\alpha, \beta=0}^3 c_{\alpha\beta} \sigma_i^{\alpha} \otimes \sigma_j^{\beta}$. The coefficients of the expansion $c_{\alpha\beta}$ are the connected correlators of two Wilson lines. The two-particle reduced density matrix is

useful in calculating the amount of entanglement *localisable* between the two chosen spins [29]. Similar results hold for three-particle (as in the GHZ-state discussed by Arvind) and, in general, for n -particle entanglement. In each of these cases, the density matrix is expressed in terms of correlators of gauge invariant observables in a topological field theory *i.e.* topological invariants [30]. The difference between two choices of the path Γ is a measure of the frustrations enclosed by the loop formed by the two paths. There is another way in which a closed loop can be obtained from an open path namely, by imposing periodic boundary conditions. If we choose this option, c_α and $c_{\alpha\beta}$ are just the expectation values of Wilson loop observables in the lattice Chern-Simons theory.

Although the Wilson loop observables are gauge-invariant, they are by no means the only interesting observables. Notice that the path connecting two points on the primary lattice pierces one plaquette on the dual lattice with every step it advances, accumulating one unit of flux in the process. Let us therefore consider the gauge-invariant operator $C = V^{-1}UV$ obtained by dressing (conjugating) the string U_{ij} by the group-valued fields V on the dual lattice. Labelling the dual lattice sites by barred coordinates, if V runs from site $\bar{i}$ to site $\bar{j}$, V^{-1} runs in the opposite direction circumnavigating the link U . We can use this operator instead of the Wilson loop to define the reduced density matrices. It may be mentioned that because of duality, the above conjugation operation simply corresponds to local unitary transformations of nearest neighbour spins on the dual lattice. Physically the operator C represents a tube of dual plaquettes whose axis lies on the primary lattice with fixed end-points.

So far there is nothing quantum about the discussion of entanglement. Indeed, the Ising spins we have considered take values ± 1 , much like classical bits; they are not allowed to be in any superposition state. The density matrices that we considered are purely thermal in nature, obtained, as they are, from the Gibbs' measure. This kind of entanglement is called thermal entanglement [31]. By using the standard Suzuki-Trotter [25] method, however, we can map the three-dimensional Ising spin glass to a two-dimensional Ising spin glass in a transverse magnetic field. The presence of the transverse magnetic field allows for transitions between the two classical states of the Ising spins and makes the system quantum mechanical. Care must be exercised in defining the spin-flip operation: σ_z and σ_x behave differently under gauge transformations because of their noncommutativity [32]. Furthermore, the situation here is slightly more complicated because the Suzuki-Trotter mapping from a d -dimensional classical statistical mechanical system to the $(d-1)$ -dimensional quantum system, requires the classical system to have different couplings along the missing (replica) dimension and the remaining dimensions. This makes the duality transformation technically a little more

involved. The results in so far as the entanglement (now truly quantum in nature) are concerned follow the same pattern. The correlation functions that appear in this case are those of the quantum spins on a two-dimensional square sub-lattice of the original three-dimensional lattice whose third dimension acts as the discretised time direction. To summarise, the reduced density matrices that one is interested in, both for thermal and quantum entanglement, are expressed in terms of expectation values of gauge invariant operators in a topological gauge theory. They are therefore topological invariants.

It is difficult to obtain a ready physical insight into the correlation functions that appear above. Let us therefore consider a simpler case. Recall that similar results hold for all abelian groups Z_p . In particular for $p \rightarrow \infty$, we have a $U(1)$ Chern-Simons gauge theory on the lattice and all the paths are continuous. In this case, it is well-known that the topological invariant produced by a computation of the correlation function of a pair of Wilson loops is the Gauss's linking number. The operator C we introduced, is the lattice generalisation of the operator introduced in the continuum BF theory by Cattaneo *et al.* [33], and gives it a nice physical interpretation as a tube of dual plaquettes. Its correlation function gives the Alexander-Conway polynomial of the (possibly) knotted axis of a plaquette-tube, with fixed end-points on the primary lattice. The partition function Z is also a topological invariant, namely, the Reidemeister torsion of the three manifold defined by the boundary conditions we choose to put on our lattice. It is equivalent to the corresponding invariant in the continuum theory, namely, the Ray-Singer analytic torsion [18]. Curiously enough, the entanglement properties of the GHZ-state which has a non-zero tripartite entanglement, but a zero bipartite entanglement cannot be accounted for, by the simple linking number invariants. This property is exactly like the corresponding property of Borromean rings which, however, are known to be distinguished from a disjoint union of unlinked rings by a higher order topological invariant, namely the Massey triple product [34]. An abelian topological theory cannot produce this invariant. It is therefore necessary to treat the spins as genuinely non-abelian objects, like one is forced to in the presence of a transverse magnetic field. In view of these remarks, the intriguing parallels between the entanglement of quantum states and the entanglement of braids, knots and links, that first appeared in [7, 9], are more realistic – may be even natural and deep-rooted – than hitherto expected.

Intuitively, topological invariants are insensitive to the presence, or changes, in length scales; this being the only distinction between a lattice and the continuum, we could borrow the relevant topological invariants from the simple continuum $U(1)$ theory in the above discussion. Such a luxury is lost if the group under consideration is finite, *e.g.* Z_2 in the case of the Ising model. Unlike

classical gauge symmetries which come from continuous local invariances, finite groups usually appear as remnants of a continuous symmetry group which is spontaneously broken. In such instances, these groups are known to give rise to cocycles in quantum field theory. This is a reflection of nontrivial group cohomology. In the present case, the finite group appears because we are dealing with spin systems. The construction of topological theories with finite groups follows from a deep result due to Dijkgraaf and Witten [13], who showed that the Chern-Simons actions for a finite group H are in one to one correspondence with the elements of the cohomology group $H^4(BH, Z)$, BH being the classifying space of H . The isomorphism $H^4(BH, Z) \approx H^3(H, U(1))$ further implies that these actions are algebraic 3-cocycles, $\omega \in H^3(H, U(1))$, which take values in $U(1)$. The presence of non-trivial group cohomology in general leads to non-trivial G bundles and the path integral involves a summation over all possible bundles. Unlike in the $U(1)$ theory, in which a determinantal expression can be derived for the partition function [35], the resulting invariants in the case of finite groups – called the Dijkgraaf-Witten invariants – are not expressible in terms of determinants. Analogues of linking numbers in topological gauge theories with finite gauge groups have also been worked out by Ferguson [36].

As the first closing remark, we mention that the details omitted in this letter can be found in a longer publication under preparation. Second, we wish to point out that many interesting problems remain. The connections between quantum entanglement and more sophisticated topological invariants like the Jones polynomial, require a nonabelian generalization of the results of this letter. Similar investigations in two and four dimensions should produce interesting connections between quantum entanglement and other important mathematical results in low dimensional topology like the intersection theory on the moduli space of Riemann surfaces, and Donaldson's invariants respectively. Finally, it is not an exaggeration to say that this letter offers a mere glimpse of a new vista which is beginning to unfold, on the relevance of finite group cohomology in the studies of entanglement in spin glasses. We hope to dilate on these issues, in the near future.

I thank M. Panero and Siddhartha Sen for some incisive comments and also A. P. Balachandran, T. R. Govindarajan, I. Tsutsui, and A. Wipf for their interest and discussions. It is a pleasure to acknowledge the hospitality of D. O'Connor and T. Dorlas of the Dublin Institute for Advanced Studies where this work was completed.

[1] L. O'Raifeartaigh *The Dawning of Gauge Theory*, (Princeton University Press, 1997).

- [2] M. A. Nielsen and I. L. Chuang *Quantum Computation and Quantum Information*, (Cambridge University Press, 2001).
- [3] F. Wilczek *Fractional Statistics and Anyon Superconductivity*, (World Scientific Publishing Company, 1990); G. Marmo, B. S. Skagerstam, A. Stern and A. P. Balachandran *Classical Topology and Quantum States*, (World Scientific Publishing Company, 1991).
- [4] J. M. Leinaas and J. Myrheim, *Nuovo Cimento* **37B**, 1 (1977).
- [5] R. Jackiw *Topological Investigations of Quantized Gauge Theories*, (Les Houches Summer School on Theoretical Physics: Relativity, Groups, and Topology, 1983); R. Rajaraman *Solitons and Instantons*, (North-Holland, 1987).
- [6] D. J. Thouless *Topological Quantum Numbers in Nonrelativistic Physics*, (World Scientific Publishing Company, 1998); M. I. Monastyrsky *Topology in Condensed Matter*, (Springer Series in Solid-State Sciences, 2005).
- [7] P. K. Arvind R. S. Cohen et al (eds), *Potentiality, Entanglement and Passion-at-a-Distance*, 53-59 (Kluwer Academic Publishers, 1997).
- [8] D. M. Greenberger, M. A. Horne and A. Zeilinger *M. Kafatos (ed), Bell's Theorem, Quantum Theory, and Conceptions of the Universe* (Kluwer Academic Publishers, 1989).
- [9] L. H. Kauffman and S. J. Lomonaco Jr., *New Journal of Physics* **4** 73, (2002).
- [10] E. Witten, *Commun. Math. Phys.* **121** 351, (1989).
- [11] J. von Neumann *Mathematical Foundations of Quantum Mechanics*, (Princeton University Press, 1996).
- [12] M. Atiyah *The Geometry and Physics of Knots*, (Cambridge University Press, 1990); E. Guadagnini *The Link Invariants of Chern-Simons Field Theory*, (Walter de Gruyter, 1993); R. K. Kaul, T. R. Govindarajan, and P. Ramadevi, hep-th/0504100 *To appear in Encyclopedia of Mathematical Physics*, Elsevier.
- [13] R. Dijkgraaf and E. Witten, *Commun. Math. Phys.* **129** 393, (1990).
- [14] *This seemingly innocuous generalisation has non-trivial repercussions. The competition between the two types of bonds leads to configurations which are frustrated. The frustrations are represented by elemental plaquettes for which the product of all the bonds is -1 . They act like dislocations or points of curvature in an otherwise flat lattice and present topological obstructions to shrinking Wilson loops to points, thus producing a Z_2 -gauge Aharonov-Bohm effect. For details, see E. Fradkin, B. A. Hluberman and S. H. Shenker, Phys. Rev. **B18** 4789-4814, (1978).*
- [15] D. Chowdhury *Spin Glasses and Other Frustrated Systems*, (World Scientific Publishing Company, 1986).
- [16] T. Senthil and M. P.A. Fisher, *Phys. Rev.* **B63** 134510, (2001).
- [17] R. Savit, *Rev. Mod. Phys.* **52** 453-487, (1980).
- [18] D. H. Adams, *Phys. Rev. Lett.* **78** 4155-4158, (1997).
- [19] S. Deser, R. Jackiw, and S. Templeton, *Phys. Rev. Lett.* **48** 975-978, (1982); *Annals of Physics* **140** 372-411, (1982).
- [20] F. Wegner, *J. Math. Phys.* **12** 2259-2272, (1971).
- [21] Samik Sen, Siddhartha Sen, J. C. Sexton, D. H. Adams, *Phys. Rev.* **E61** 3174-3185, (2000).
- [22] R. Kantor and L. Susskind, *Nuc. Phys.* **B366** 533-568, (1991).
- [23] A. Yu Kitaev, *Annals of Physics* **303** 2-30, (2003); M. H. Freedman, A. Kitaev, M. J. Larsen, and Z.

- Wang, Bull. Am. Math. Soc. **40** No. 1, 31-38, (2002); J. Preskill, Caltech Ph219 Lecture Notes, Chapter 9, <http://www.theory.caltech.edu/preskill/ph229>
- [24] I. Chiorescu, Y. Nakamura, C. J. P. M. Harmans, J. E. Mooij, Science **299** 1869, (2003); I. Chiorescu, P. Bertet, K. Semba, Y. Nakamura, C.J.P.M. Harmans, J. E. Mooij, Nature **431** 159, (2004).
- [25] H. Nishimori *Statistical Physics of Spin Glasses and Information Processing: An Introduction*, (Oxford University Press, 2001).
- [26] N. Surlas, Nature **339** 693, (1989).
- [27] X. G. Wen and A. Zee, Phys. Rev. **B46** 2290-2301, (1992).
- [28] D. Birmingham, M. Blau, M. Rakowski, and G. Thompson, Phys. Rep. **209**, 129-340, (1991).
- [29] F. Verstraete, M. Popp and J. I. Cirac, Phys. Rev. Lett. **92** No.2 027901, (2004).
- [30] *It is in general true that the reduced density matrices are completely determined by the correlation functions of physical observables; these, however, are in general difficult to calculate exactly. For a representative list see*, T. J. Osborne and M. A. Nielsen, Phys. Rev. **A66** 032110, (2002); S. Bose, Phys. Rev. Lett. **91** 207901, (2003); A. Lakshminarayan and V. Subrahmanyam, Phys. Rev. **A67** 052304, (2003); V. Subrahmanyam, Phys. Rev. **A69** 022311, (2004); Phys. Rev. **A69** 034304, (2004).
- [31] *In the field of critical phenomena, it has taken a curiously long time to coin the name quantum phase transitions for macroscopic changes of state driven by parameters other than temperature e.g. magnetic fields. In contrast, the realisation that entanglement between spins can be established by purely thermal - as opposed to quantum correlations - has dawned fairly early. See, for example* M. C. Arnesen, S. Bose and V. Vedral, Phys. Rev. Lett. **87** No.1 017901, (2001).
- [32] S. Morita, Y. Ozeki, H. Nishimori, J. Phys. Soc. of Japan **75** No. 1 014001, (2006).
- [33] A. S. Cattaneo, P. Cotta-Ramusino, M. Martellini, Nuc. Phys. **B436**, 355-384, (1995).
- [34] W. S. Massey, Journal of Knot Theory and Its Ramifications, **7** No. 3, 393-414, (1998).
- [35] A. S. Schwarz, Lett. Math. Phys. **2** 247-252, (1978).
- [36] Kenneth Ferguson, Journal of Knot Theory and Its Ramifications, **2** No. 1, 11-36, (1993).